



72569754231 294640481.83333 5903142.7340426 32882497388 9171979594 94935093120 36225918.37931 20104449.089744 177018732016 951617664.5 63226570930 92289036840 14894060.228571 133278879396 23961615.235294

Hyperemesis gravidarum acog pdf online for pc

Acog hyperemesis gravidarum. Hyperemesis acog.

Journal Of Maternal-Fetal & Amp; Neonatal Medicine: The Official Journal Of The European Association Of Perinatal Medicine, The Federation Of Asia And Oceania Perinatal Society of Perinatal Obstetricians 22(4):353 [PubMed] [Google Scholar]Trogstad LIS, Stoltenberg C, Magnus P, Skjaerven R, Irgens LM (2005) Recurrence risk in hyperemesis gravidarum. Arch Gynecol Obstet 268(3):219. A marginally higher rate of birth defects was observed in the mirtazapine and SSRI groups compared with the low rate of birth defects in the general control subjects, which is consistent with other studies that evaluated the risk factors related to the underlying depressive disorder (Huybrechts et al. [PubMed] [Google Scholar]Seto A, Einarson T, Koren G (1997) Pregnancy outcome following first trimester exposure to antihistamines: meta-analysis. 2013).Dopamine antagonists, such as metoclopramide and promethazine and droperidol, have been used to treat HG (Fadi et al. Female fetuses are more commonly born to women with HG (Veenendaal et al. Midwifery 25(6):649. J Clin Psychopharmacol 35(3):250. Am J Obstet Gynaecol 31(8):708. Am J Perinatol 15(9):533. 1997; Tan et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et al. Both medications resulted in similar improvement in N/V). al. 2000) (Gadsby et al. However, there are several case studies reporting the efficacy of hypnotherapy (Fuchs et al. 2000). Am J Perinatol 16(2):85. Rates of mirtazapine for HG to summarize the published clinical data. Am J Obstet Gynecol 186(5):S228. The Journal of Midwifery & Women's Health 56(2):132-136 [PMC free article] [PubMed] [Google Scholar] Fejzo MS, Magtira A, Schoenberg FP, Macgibbon K, Mullin PM (2015) Neurodevelopmental delay in children exposed in utero to hyperemesis gravidarum. However, other studies have found that ondansetron was associated with birth defects. Obstetrics & Gynecology 102(6):1250. [PubMed] [Google Scholar]Simon EP, Schwartz J (1999) Medical Hypnosis for Hyperemesis Gravidarum Birth 26(4):248. Am Fam Physician 81(9):1139-1142 [PubMed] [Google Scholar]ACOG (2015) Practice bulletin 153: nausea and vomiting of pregnancy. 1959; Koren et al. [PubMed] [Google Scholar]Kashifard M, Basirat Z, Kashifard M, Golsorkhtabar-Amiri M, Moghaddamnia A (2013) Ondansetrone or metoclopromide? 2011; Sheehan 2007; Simpson et al. Acta Obstet Gynecol Scand 94(4):359–367 [PubMed] [Google Scholar]Summers A (2012) Emergency management of hyperemesis gravidarum: Anthonysummersdiscussesthecauses, signs and symptoms of severe morning sickness, and explains how the condition should be treated in emergency care settings. [PubMed] [Google Scholar]Kutcher JS, Engle A, Firth J, Lamm SH (2003) Bendectin and birth defects. [PubMed] [Google Scholar]McKeigue P, Lamm S, Linn S, Kutcher J (1994) Bendectin and birth-defects. [PubMed] [Google Scholar]Kutcher JS, Engle A, Firth J, Lamm SH (2003) Bendectin and birth-defects. Additionally, a second Danish study using the same registry but including more infants over more years found a two-fold increase in cardiac malformations (Andersen et al. Aust Fam Physician 36(9):698. Autonomic Neuroscience: Basic And Clinical 129(1):22. 2007). 2011) (Fejzo et al. [PubMed] [Google Scholar]Emelianova S, Mazzotta P, Einarson A, Koren G (1999) Prevalence and severity of nausea and vomiting of pregnancy and effect of vitamin supplementation. [PubMed] [Google Scholar]Kimura M, Amino N, Tamaki H, Ito E, Mitsuda N, Miyai K, Tanizawa O (1993) Gestational thyrotoxicosis and hyperemesis gravidarum: possible role of hCG with higher stimulating activity. Drug Saf 19(2):155. 1996). A case report. For the majority of women, N/V resolves by the 16th week of pregnancy (Eliakim et al. [PubMed] [Google Scholar]Rhodes VA, McDaniel RW (1999) The index of nausea, vomiting, and retching: a new format of the index of nausea and vomiting. (Clinical report). 2003; Van Thiel et al. Am J Obstet Gynecol 186(5):S253. Am J Epidemiol 149(8):717. 1993). [PubMed] [Google Scholar]Jamigorn M, Phupong V (2007) Acupressure and vitamin B6 to relieve nausea and vomiting in pregnancy: a randomized study. Acta Obstet Gynecol Scand 88(11):1190. Journal Of The Medical Association Of Thailand = Chotmaihet Thangphaet 90(1):15. 2002; Seto et al. Obstet Gynecol 126:687-688 [PubMed] [Google Scholar]Aghadam SKZ, Mahfoozi B (2010) Evaluation of the effects of acupressure by sea band on nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Aghadam SKZ, Mahfoozi B (2010) Evaluation of the effects of acupressure by sea band on nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Aghadam SKZ, Mahfoozi B (2010) Evaluation of the effects of acupressure by sea band on nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Aghadam SKZ, Mahfoozi B (2010) Evaluation of the effects of acupressure by sea band on nausea and vomiting of pregnancy. Liberzon I (2007) Service use data analysis of pre-pregnancy psychiatric and somatic diagnoses in women with hyperemesis gravidarum. Acupressure treatment of morning sickness in pregnancy. Reprod Toxicol 9(4):337. Birth defects research. [PubMed] [Google Scholar]Yost NP, McIntire DD, Wians FH, Ramin SM, Balko JA, Leveno KJ (2003) A randomized, placebo-controlled trial of corticosteroids for hyperemesis due to pregnancy. PLoS One 9(3) [PMC free article] [PubMed] [Google Scholar]Tasci Y, Demir B, Dilbaz S, Haberal A (2009) Use of diazepam for hyperemesis gravidarum. Psychiatry Clin Neurosci 62(1):75. International Journal Of Women&Apos;S Health 6:719 [PMC free article] [PubMed] [Google Scholar]McCormack D (2010) Hypnosis for hyperemesis gravidarum. Obstetrics & Gynecology 100(1):94. 2006). 2003; McKeigue et al. 2015) conducted a prospective cohort study comparing birth defects in infants of women treated antenatally with mirtazapine to two comparison groups, each with 357 subjects. Innovations In Clinical Neuroscience 9(11-12):31 [PMC free article] [PubMed] [Google Scholar]Sripramote M, Lekhyananda N (2003) A randomized comparison of ginger and vitamin B6 in the treatment of nausea and vomiting of pregnancy. Obstetrics & Gynecology 91(1):78. 2009), although another larger study of 8-year-old children of mothers with HG, regardless of treatment received, showed significantly increased rates of neurodevelopmental delay (Fejzo et al. 1992). [PubMed] [Google Scholar]Gideon K, Katayoon M, Caroline M, Simerpal Kaur G (2009) The effect of acid-reducing pharmacotherapy on the severity of nausea and vomiting of pregnancy. Am J Perinatol 14(3):119. [PubMed] [Google Scholar]Freedman SB, Uleryk E, Rumantir M, Finkelstein Y (2013) Ondansetron and the risk of cardiac arrhythmias: a systematic review and postmarketing analysis. [PubMed] [Google Scholar]Pasricha PJ, Nonko P, Aravind S, Joseph J (2006) Drug insight: from disturbed motility to disordered movement—a review of the clinical benefits and medicolegal risks of metoclopramide. 2000), between 15 and 20% of women continue to experience symptoms until the third trimester and 5% through delivery (Goodwin 2008) (Fell et al. [PubMed] [Google Scholar]Geiger CJ, Fahrenbach DM, Healey FJ (1959) Bendectin in the treatment of nausea and vomiting in pregnancy. Seven patients presented with symptoms of depression and five reported a history of depression. [PubMed] [Google Scholar]Wallstedt A, Riely CA, Shaver D (1990) Prevalence and characteristics of liver dysfunction in hyperemesis gravidarum. Women presenting with HG are evaluated with serial measures of maternal weight, orthostatic blood pressures, heart rate, electrolytes, urine ketones, and an obstetric ultrasound if not previously obtained (2015). Journal Of Alternative And Complementary Medicine (New York, NY) 15(3):243 [PubMed] [Google Scholar]Park-Wyllie L, Mazzotta P, Pastuszak A, Moretti ME, Beique L, Hunnisett L, Friesen MH, Jacobson S, Kasapinovic S, Chang D, Diav-Citrin O, Chitayat D, Dia Nulman I, Einarson TR, Koren G (2000) Birth defects after maternal exposure to corticosteroids: prospective cohort study and meta-analysis of epidemiological studies. Acad Emerg Med 15(3):209 [PubMed] [Google Scholar]Brent RL (1995) Bendectin: review of the medical literature of a comprehensively studied human nonteratogen and the most prevalent tortogen-litigen. 2014; Pasricha et al. [PubMed] [Google Scholar]Morali JA, Braverman DZ (1990) Abnormal liver enzymes and ketonuria in hyperemesis gravidarum: a retrospective review of eighty patients. 2007). Mirtazapine acts on noradrenergic, serotonergic, histaminergic, and muscarinic receptors. Pharmacoepidemiol Drug Saf 22:13 [Google Scholar] Annagür BB, Kerimoğlu ÖS, Gündüz Ş, Tazegül A (2014) Are there any differences in psychiatric symptoms and eating attitudes between pregnant women with hyperemesis gravidarum and healthy pregnant women? 2000) before starting mirtazapine and educated about the rare possibility of anxiety, agitation, and suicidal ideation at the beginning of treatment. Mirtazpine is metabolized in the liver and its elimination half-life is between 20 and 40 h. [PubMed] [Google Scholar]Smith C, Crowther C, Willson K, Hotham N, McMillian V (2004) A randomized controlled trial of ginger to treat nausea and vomiting in pregnancy. In 2008, the rate of prescriptions per month. Clin Endocrinol 38(4):345 [PubMed] [Google Scholar]Klebanoff MA, Koslowe PA, Kaslow R, Rhodes GG (1985) Epidemiology of vomiting in early pregnancy. Arch Gynecol Obstet 276(3):245. [PubMed] [Google Scholar]Spiegel DR, Webb K (2012) A case of treatment refractory hyperemesis gravidarum in a patient with comorbid anxiety, treated successfully with adjunctive gabapentin: a review and the potential role of neurogastroentereology in understanding its pathogenesis and treatment. 1994; Neutel and Johansen 1995). Am J Obstet Gynecol 186(5):S256. N Engl J Med 371 (12): 1168-1169 [PubMed] [Google Scholar]Jackson CW, Sheehan AH, Reddan JG (2007) Evidence-based review of the black-box warning for droperidol.(SPECIAL FEATURE)(drug overview)(clinical report). [PubMed] [Google Scholar]Jackson CW, Sheehan AH, Reddan JG (2007) Evidence-based review of the black-box warning for droperidol.(SPECIAL FEATURE)(drug overview)(clinical report). [PubMed] [Google Scholar]Jackson CW, Sheehan AH, Reddan JG (2007) Evidence-based review of the black-box warning for droperidol.(SPECIAL FEATURE)(drug overview)(clinical report). [PubMed] [Google Scholar]Jednak MA, Shadigian EM, Kim MS, Woods ML, Hooper FG, Owyang C, Hasler WL (1999) Protein meals reduce nausea and gastric slow wave dysrhythmic activityinfirst trimester pregnancy. 2008). J Clin Endocrinol Metab 81:1160-1163 [PubMed] [Google Scholar]Poursharif B, Korst LM, Fejzo MS, Macgibbon KW, Romero R, Goodwin TM (2007) The psychosocial burden of hyperemesis gravidarum. Journal Of The Medical Association Of Thailand = Chotmaihet Thangphaet 86(9):846. Oncol Nurs Forum 26(5):889. [PubMed] [Google Scholar]Koren G, Clark S, Hankins GDV, Caritis SN, Miodovnik M, Umans JG, Mattison DR (2010) Effectiveness of delayed-release doxylamine and pyridoxine for nausea and vomiting of pregnancy: a randomized placebo controlled trial. Obstet Gynecol 123(6):1272. 2002; Maltepe and Koren 2013b; Wheatley 1977) and safety (Brent 1995; Einarson et al. [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. J Clin Psychopharmacol 34(4):540-541 [PubMed] [Google Scholar]Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are Scholar]Anttila SAK, Leinonen EVJ (2001) A review of the pharmacological and clinical profile of mirtazapine. [PubMed] [Google Scholar]Bischoff SC, Renzer C (2006) Comparing ginger and vitamin B6 for the treatment of nausea and vomiting in pregnancy: a randomized controlled trial. [PubMed] [Google Scholar]Bischoff SC, Renzer C (2006) Nausea and nutrition. [PubMed] [Google Scholar]Zhang Y, Cantor RM, MacGibbon K, Romero R, Goodwin TM, Mullin PM, Fejzo MS (2011) Familial aggregation of hyperemesis gravidarum. [PubMed] [Google Scholar]Ansermot N, Hodel PF, Eap CB (2014) Serotonin toxicity after addition of mirtazapine to escitalopram. Electrolyte abnormalities are corrected with intravenous replacement fluids; however, in severe cases, sustained weight loss requires enteral or total parenteral nutrition (Ayyavoo et al. Metoclopramide was found to be more effective at reducing N/V than other dopamine antagonists. Am J Obstet Gynecol 211(2) [PubMed] [Google Scholar]Norheim AJ, Pedersen EJ, Fønnebø V, Berge L, 2001. Bjog : An International Journal Of Obstetrics And Gynaecology 112(12):1641. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Källén B, Lundberg A (2003) Relationship between vitamin use, smoking A (2003) Relationship betwee abnormal in patients with severe hyperemesis gravidarum. 2014), and a US cohort found an increased risk of cleft palate (Anderka et al. Side effects include headache; fatigue; constipation; and, rarely, serotonin syndrome (Freedman et al. J Clin Endocrinol Metab 75(5):1333. The comparison groups were (1) mothers treated with selective serotonin reuptake inhibitor antidepressants and (2) and a general population of women without antidepressant treatment or exposure to known teratogenic agents. 2005). Obstet Gynecol Clin N Am 35(3):401 [PubMed] [Google Scholar]Goodwin TM, Hershman JM, Cole L (1994) Increased concentration of the free beta-subunit of human chorionic gonadotropin in hyperemesis gravidarum. Mirtazapine has been associated with serotonin syndrome in multiple case reports, typically when multiple serotonergic drugs are combined (Ansermot et al. European Journal of Obstetrics and Gynecology 189:79 [PubMed] [Google Scholar]Fell DB, Dodds L, Joseph KS, Allen VM, Butler B (2006) Risk factors for hyperemesis gravidarum requiring hospital admission during pregnancy. [PubMed] [Google Scholar]Rohde A, Dembinski J, Dorn C (2003) Mirtazapine (Remergil) for treatment resistant hyperemesis gravidarum: rescue of a twin pregnancy. Obstet Gynecol 78(1):33. Am J Obstet Gynecol 186(5):S210. 2014; Tamay and Kuscu 2011).Antihistamines (H1 antagonists), including doxylamine used in combination with pyridoxine, meclizine, dimenhydrinate, and diphenhydramine, are effective in reducing N/V. J Perinatol 28(3):176-181 [PubMed] [Google Scholar]Quigley EMM, Hasler WL, Parkman HP (2001) AGA technical review on nausea and vomiting. Am J Obstet Gynecol 210(6):521. Am J Obstet Gynecol 173(3):881. European Journal Of Obstetrics And Gynecology 141(1):13 [PMC free article] [PubMed] [Google Scholar]Fejzo MS, MacGibbon KW, Romero R, Goodwin TM, Mullin PM(2011) Recurrence risk of fetal malformations (Matok et al. [PubMed] [Google Scholar]Li L, Li L, Zhou X, Xiao S, Gu H, Zhang G, 2015. As progesterone increases throughout gestation, it is associated with decreased esophageal and gastric motility as well as lower esophageal sphincter pressure. [PubMed] [Google Scholar]Saks BR (2001) Mirtazapine: treatment of depression, anxiety, and hyperemesis gravidarum in the pregnant patient. Although it usually occurs between weeks 4 and 9 of pregnancy and resolves by mid-gestation (Eliakim et al. Am J Physiol 277(4 Pt 1):G855. 2010a). 2006).Women with HG often present with increased blood urea nitrogen and hematocrit and, in 15-25% of cases, hyponatremia, hypokalemia, and hypochloremia (Goodwin 1998). The options are limited for women who are not adequately treated with these medications. Also, metoclopramide with pyridoxine was also more effective than were other dopamine antagonists, prochlorperazine and promethazine (Fadi et al. HG is the most common indication for hospital admission in the first 20 weeks of pregnancy (Klebanoff et al. Obstet Gynecol 72(1):102. Obstetrics & Gynecology 101(4):639. Ondansetron is effective in reducing N/V during pregnancy, and a double-blind randomized controlled study of 36 women found that ondansetron was significantly more effective than combined pyridoxine and doxylamine treatment in reducing nausea and vomiting (Oliveira et al. Obstetrics & Gynecology 88(3):343. Obstet Gynecol 107(2 Pt 1):277. 2013). [PubMed] [Google Scholar]Vikanes Å, Grjibovski ÅM, Vangen S, Gunnes N, Samuelsen SO, Magnus P (2010) Maternal body composition, smoking, and hyperemesis gravidarum. 2003; Saks 2001; Schwarzer et al. 1980; McCormack 2010; Simon and Schwartz 1999). Acta Neurol Belg 93:276-282 [PubMed] [Google Scholar]Pradat P, Robert-Gnansia E, Di Tanna GL, Rosano A, Lisi A, Mastroiacovo P (2003) First trimester exposure to corticosteroids and oral clefts. [PMC free article] [PubMed] [Google Scholar] Gastroenterology 120(1):263. 1990). HG is a clinical diagnosis; however, tools such as the Pregnancy Unique Quantification of Emesis (PUQE) score or Rhodes index have been used to assess the severity of symptoms in research studies (Koren et al. 1992; Emelianova et al. Clin Obstet Gynecol 41: 597-605 [PubMed] [Google Scholar]Goodwin TM (2008) Hyperemesis Gravidarum. [PubMed] [Google Scholar]Stokke G, Gjelsvik BL, Flaatten KT, Birkeland E, Flaatten H, Trovik J (2015) Hyperemesis gravidarum, nutritional treatment by nasogastric tube feeding: a 10-year retrospective cohort study. [PubMed] [Google Scholar]Gadsby R, Barnie-Adshead AM, Jagger C (1993) A prospective study of nausea and vomiting during pregnancy. (Understanding and treating nausea and vomiting of pregnancy). [PubMed] [Google Scholar]Fitzgerald JP (1955) The effect of promethazine in nausea and vomiting of pregnancy. [PMC free article] [PubMed] [Google Scholar]Liu M-C,Kuo S-H,Lin C-P,Yang Y-H(2014)Effects of professional support on nausea, vomiting, and quality of life during early pregnancy. Int J Clin Exp Hypn 28(4):313. Am J Perinatol 17(4):207. 2013). The safety of ondansetron during pregnancy was assessed in a Danish study of 1970 exposed infants, which did not show an increased risk of fetal malformations or adverse pregnancy outcomes (Pasternak et al. [PubMed] [Google Scholar]Matthews A, Haas DM, Mathúna DPO, Dowswell T, Doyle M (2014) Interventions for nausea and vomiting in early pregnancy. 1985) (Depue et al. Such studies should determine characteristics of women who would benefit most from each treatment option to personalize care. Abas MN, Tan PC, Azmi N, Omar SZ (2014) Ondansetron compared with metoclopramide for hyperemesis gravidarum: a randomized controlled trial. [PubMed] [Google Scholar]Huxley RR (2000) Nausea and vomiting in early pregnancy: its role in placental development. European Journal Of Obstetrics And Gynecology 143(1):43 [PubMed] [Google Scholar]Lagiou P, Tamimi R, Mucci LA, Trichopoulos D, Adami H-O, Hsieh C-C (2003) Nausea and vomiting in pregnancy in relation to prolactin, estrogens, and progesterone: a prospective study. [PubMed] [Google Scholar]Simpson SW, Goodwin TM, Robins SB, Rizzo AA, Howes RA, Buckwalter JG (2001) Psychological factors and hyperemesis gravidarum. BJOG: An International Journal of Obstetrics and Gynaecology 121(12):1556. Am J Health Syst Pharm 64(11):1174. Obstet Gynecol 115(5):975. Primiparous women; adolescents; women with increased BMI, multiple gestations, and gestational trophoblastic disease; and women with HG during a previous pregnancy also have increased risk for HG (Klebanoff et al. 2003). Hormonal factors have been evaluated as etiologic contributors to HG. 2014). Historically, HG was thought to be a response to stress or ambivalence about pregnancy (Buckwalter and Simpson 2002); however, no differences in psychological or personality characteristics between women with and without HG have been demonstrated (Orazio et al. It is an inverse agonist on the H1 receptor which contributes to the sedative effect. [PubMed] [Google Scholar]Huybrechts KF, Hernández-Díaz S, Avorn J (2014) Antidepressant use in pregnancy and the risk of cardiac defects. Journal of Women's Health 20(1):137 [PMC free article] [PubMed] [Google Scholar]Ozgoli G, Goli M, Simbar M (2009) Effects of ginger capsules on pregnancy, nausea, and vomiting. 2015; Liu et al. Arch Gynecol Obstet 291(6):1271. Clinical and investigative medicine. 2015). There are few studies examining psychotherapy treatment for HG, including no randomized trials (2015). Journal Of Obstetrics And Gynaecology 30(7):647. Am J Med Genet 86(3):242. It is an indirect agonist of the 5-HT1A receptor, an antagonist at the 5-HT2 receptors, and an inverse agonist at the 5-HT2C receptor to produce the antidepressant effect (Blier and Abbott 2001). [PubMed] [Google Scholar]Chittumma P, Kaewkiattikun K, Wiriyasiriwach B (2007) Comparison of the effectiveness of ginger and vitamin B6 for treatment of nausea and vomiting in early pregnancy: a randomized double-blind controlled trial. [PubMed] [Google Scholar]Maltepe C, Koren G, 2013a. Iranian Journal of Obstetrics, Gynecology and Infertility 13(2):39 [Google Scholar]Anderka M, Mitchell AA, Louik C, Werler MM, Hernández-Diaz S, Rasmussen SA (2012) Medications used to treat nausea and vomiting of pregnancy and the risk of selected birth defects. No statistically significant difference in the rate of major birth defects after first-trimester exposure between mirtazapine, SSRI-exposed, and non-exposed pregnancies was found. When metoclopramide was compared to ondansetron, they were comparable in reducing nausea, although ondansetron had a greater effect on reducing vomiting (Kashifard et al. [PubMed] [Google Scholar]Hallak M, Tsalamandris K, Drombrowski MP, Isada NB, Pryde PG, Evans MI (1996) Hyperemesis gravidarum: effects on fetal outcome. Women with a sister or mother who had HG are also at increased risk, which suggests that genetic factors play a role (Zhang et al. Which is more effective in severe nausea and vomiting of pregnancy? [PubMed] [Google Scholar]Tan PC, Vani S, Lim BK, Omar SZ (2010b) Anxiety and depression in hyperemesis gravidarum: prevalence, risk factors and correlation with clinical severity. Diclegis® is considered first-line pharmacotherapy for N/V in pregnancy by the American Professors in Gynecology and Obstetrics and the American College of Obstetricians and Gynecologists (2015). II: ecological analyses. [PubMed] [Google Scholar]Shepard TH, Brent RL, Friedman JM, Jones KL, Miller RK, Moore CA, Polifka JE (2002) Update on new developments in the study of human teratogens. 2006; Trogstad et al. Am J Obstet Gynecol 193(3):811. Biological Research For Nursing 16(4):378. Similar to other antidepressants, patients should under-go screening for bipolar disorder using an instrument such as the Mood Disorder Questionnaire (Hirschfeld et al. Mirtazapine may expand the options for treating HG, especially in women who are also experiencing psychiatric symptoms such as depression and anxiety. [PubMed] [Google Scholar]Murphy PA (1998) Alternative therapies for nausea and vomiting of pregnancy. Archives Of Women's Mental Health 3(4):165 [Google Scholar]Murphy PA (1998) Alternative therapies for nausea and vomiting of pregnancy. Archives Of Women's Mental Health 3(4):165 [Google Scholar]Murphy PA (1998) Alternative therapies for nausea and vomiting of pregnancy. analysis of case-control studies. Antagonism of the 5-HT2 receptor also contributes to the anxiolytic, sedative, and appetite-stimulating effects (Fawcett and Barkin 1998). (2016) concluded that little evidence existed to support the superiority of one intervention over another in the treatment of HG, and the authors recommended research comparing the side-effect profiles, safety, economic costs, and benefits of treatments to inform selection. [PubMed] [Google Scholar]Hinkle Stefanie N., Mumford Sunni L., Grantz Katherine L., Silver Robert M., Mitchell Emily M., Sjaarda Lindsey A., Radin Rose G., Perkins Neil J., Galai Noya, Schisterman Enrique F. 1994). Several investigators have reported and increased level of free β -hCG in women with HG compared to matched controls (Goodwin et al. [PubMed] [Google Scholar]Tsang IS, Katz VL, Wells SD (1996) Maternal and fetal outcomes in hyperemesis gravidarum. Bjog-An International Journal Of Obstetrics And Gynaecology 118(11):1302. Am J Obstet Gynecol 174(6):1801. Eur J Obstet Gynecol Reprod Biol 149(2):153. A systematic review and meta-analysis of the effect and safety of ginger in the treatment of pregnancy-associated nausea and vomiting. The criteria that define HG include dehydration, ketonuria, and more than 5% weight loss (Nelson-Piercy 1998) (McCarthy et al. 2003; Vikanes et al. Reviews In Obstetrics & Amp; Gynecology 5(2):78 [PMC free article] [PubMed] [Google Scholar]Weigel RM, Weigel MM (1989) Nausea and vomiting of early-pregnancy and pregnancy and pregn Gynecol Obstet 251(4):181. [PubMed] [Google Scholar]Veenendaal M, van Abeelen A, Painter RC, van Der Post J, Roseboom T (2011) Consequences of hyperemesis gravidarum for offspring: a systematic review and meta-analysis. 1986); however, these findings were not confirmed in another investigation (Wilson et al. Additionally, hCG peaks in the first trimester when N/V is typically the most severe. [PubMed] [Google Scholar]Sw Kim, Is Shin, Jm Kim, Yc Kim, Ks Kim, Km Kim, Sj Yang, Js Yoon (2008) Effectiveness of mirtazapine for nausea and insomnia in cancer patients with depression. Am J Obstet Gynecol 160:906-909 [PubMed] [Google Scholar]Sw Kim, Is Shin, Jm Kim, Yc Kim, Ks Kim, Km Kim, Sj Yang, Js Yoon (2008) Effectiveness of mirtazapine for nausea and insomnia in cancer patients with depression. Mirtazapine use in resistant hyperemesis gravidarum: report of three cases and review of the literature. [PubMed] [Google Scholar]Nulman I, Rovet J, Barrera M, Knittel-Keren D, Feldman BM, Koren G (2009) Long-term neurodevelopment of children exposed to maternal nausea and vomiting of pregnancy and diclectin. 1999; Källén et al. Journal Of Maternal-Fetal & Amp; Neonatal Medicine: The Official Journal Of The European Association Of Perinatal Medicine, The Federation Of Asia And Oceania Perinatal Society of Perinatal Society of Perinatal Obstetricians 25(6):632 [PMC free article] [PubMed] [Google Scholar]Mullin PM, Goodwin A, Fejzo F, Bray KW, Schoenberg R, MacGibbon TM, Romero MS, Fejzo MS, 2011. 2010; Magee et al. 1985) (Price et al. Br J Obstet Gynaecol 96(11):1312. 2009; Pasternak et al. J Clin Gastroenterol 12:303-305 [PubMed] [Google Scholar]Mullin PM, Ching C, Schoenberg F, Macgibbon K, Romero R, Goodwin TM, Fejzo MS (2012) Risk factors, treatments, and outcomes associated with prolonged hyperemesis gravidarum. J Reprod Med 41:871-874 [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. [PubMed] [Google Scholar]Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea therapies for severe or refractory HG, which has been associated with requests for termination of desired pregnancies, is imperative. Diclegis® received FDA approval in 2013 with a Pregnancy C (1998) Treatment of nausea and vomiting in pregnancy. A total of 15 reports of successful treatment of HG with mirtazapine have been published. [PubMed] [Google Scholar]Shrim A, Boskovic R, Maltepe C, Navios Y, Garcia-Bournissen F, Koren G, Shrim G (2006) Pregnancy outcome following use of large doses of vitamin B 6 in the first trimester. Comorbid conditions, most commonly reflux, should also be evaluated and treated (Gideon et al. [PubMed] [Google Scholar]Davis M (2004) Nausea and vomiting of—an evidence-based review. [PubMed] [Google Scholar]Nageotte MP, Briggs GG, Towers CV, Asrat T (1996) Droperidol and diphenhydramine in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Schwarzer V, Heep A, Gembruch U, Rohde A (2008) Treatment resistant hyperemesis gravidarum in a patient with type 1 diabetes mellitus: neonatal withdrawal symptoms after successful antiemetic therapy with mirtazapine. Arch Gynecol Obstet 277(1):67. [PubMed] [Google Scholar]Buckwalter JG, Simpson SW (2002) Psychological factors in the etiology and treatment of severe nausea and vomiting in pregnancy. A severe and prolonged form of N/V, hyperemesis gravidarum (HG), affects 0.3-2% of pregnancies (Hod et al. Pak J Biol Sci 14(16):817. Mirtazapine exerts its antiemetic effect regardless of whether the patient has a psychiatric disorder, though a patient with depression or anxiety may benefit from both of these actions. Mirtazapine's side effects were rated as average compared to other antidepressants and commonly include sedation, weight gain, dry mouth, and constipation (Cipriani et al. Mirtazapine has highly promising but efficacy and side effect data are limited to multiple case studies. J Obstet Gynaecol 26(8):749. Teratology 50(1):27. Nutricion hospitalaria 21(Suppl 3):17-26 [PubMed] [Google Scholar]Carmichael SL, Shaw GM (1999) Maternal corticosteroid use and risk of selected congenital anomalies. (2016) Association of Nausea and Vomiting During Pregnancy With Pregnancy Loss. 1993). There is homology between the beta-subunit of hCG and TSH which causes hCG to have thyroidstimulating activity (Ballabio et al. Br J Obstet Gynaecol 84(6):444. 2002; Rhodes and McDaniel 1999). Emergency Nurse 20(4):24 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect. Nutrition Journal 13. Journal Of Women&Apos; S Health & Amp; Gender-Based Medicine 10(5):471 [PubMed] [Google Scholar]Tamay AG, Kuscu N (2011) Hyperemesis gravidarum: current aspect Scholar]Slaughter SR, Hearns-Stokes R, Van Der Vlugt T, Joffe HV (2014) FDA approval of doxylamine-pyridoxine therapy for use in pregnancy. N Engl J Med 370(12):1081. Hum Reprod 30(12):2764-2773 [PubMed] [Google Scholar]Fawcett J, Barkin RL (1998) Review of the results from clinical studies on the efficacy, safety and tolerability of mirtazapine for the treatment of patients with major depression. N Engl J Med 368(9):814. pylori has been shown to improve N/V in infected women (El Younis et al. J Pediatr 155(1):45. Though no case reports of serotonin syndrome have been found with the combination of mirtazapine and metoclopramide or ondansetron, it remains a theoretical risk. The risk of recurrent HG is 15-20% (Dodds et al. Initial pharmacologic treatment for HG includes a combination of doxylamine and pyridoxine. Acta Endocrinol 111:404-410 [PubMed] [Google Scholar]Boelig Rupsa C., Barton Samantha J., Saccone Gabriele, Kelly Anthony J., Edwards Steve J., and Berghella Vincenzo (2016) Interventions for treating hyperemesis gravidarum." Cochrane Database of Systematic Reviews Reviews [PubMed] [Google Scholar] Boneva RS, Moore CA, Botto L, Wong LY, Erickson JD (1999) Nausea during pregnancy and congenital heart defects: a population-based case-control study. Clinical and experimental obstetrics & gynecology 40(1):127-130 [PubMed] [Google Scholar]Keating A, Chez RA (2002) Ginger syrup as an antiemetic in early pregnancy. Obstet Gynecol Int 2009(2009) [PMC free article] [PubMed] [Google Scholar]Keating A, Chez RA (2002) Ginger syrup as an antiemetic in early pregnancy. Obstet Gynecol Int 2009(2009) [PMC free article] [PubMed] [Google Scholar]Keating A, Chez RA (2002) Ginger syrup as an antiemetic in early pregnancy. infection: a systematic review. 1985) (Eliakim et al. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google Scholar]Hsu J], Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. [PubMed] [Google S pregnancy: a randomized, double-blind placebo-controlled study. Preemptive treatment of nausea and vomiting of pregnancy: results of a randomized controlled trial. 2008; Uguz, 2014). In a recent Cochrane review, Boelig, et al. Droperidol was studied in combination with diphenhydramine and was found to reduce days in the hospital for HG (Nageotte et al. Journal Of Perinatal &Amp; Neonatal Nursing 18(4):312 [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. JAMA Inter Med [PMC free article] [PubMed] [Google Scholar]Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemion factors: a seroepidemion fa Scholar]Hirschfeld RMA, Williams JBW, Spitzer RL, Calabrese JR, Flynn L, Keck PE, Lewis L, Mcelroy SL, Post RM, Rapport DJ, Russell JM, Sachs GS, Zajecka J (2000) Development and validation of a screening instrument for bipolar spectrum disorder: the mood disorder questionnaire. [PubMed] [Google Scholar]Fejzo MS, Ingles SA, Wilson M, Wang W, MacGibbon K, Romero R, Goodwin TM (2008) High prevalence of severe nausea and vomiting of pregnancy and hyperemesis gravidarum among relatives of affected individuals. 1987). 2001). [PubMed] [Google Scholar]Danielsson B, Wikner BN, Källén B (2014) Use of ondansetron during pregnancy and congenital malformations in the infant. Mirtazapine is a weak inhibitor of CYP-isoenzymes and is unlikely to cause many clinically relevant drug interactions (Anttila and Leinonen 2001). [PubMed] [Google Scholar]Bashiri A, Newmann L, Maymon E, Katz M (1995) Hyperemesis gravidarum: epidemiologic features, complications and outcome. 2015). Ondansetron, a serotonin antagonist acting at the 5-HT3 receptor, is the most commonly prescribed medication for the treatment of N/V during pregnancy, and its use is rapidly increasing (Koren 2014). p 200. Obstet Anesth Dig 34(4):211 [Google Scholar]Peeters A, van de Wyngaert F, Van Lierde M, Sindic CJ, Laterre EC (1993) Wernicke's encephalopathy and central pontine myelinolysis induced by hyperemesis gravidarum. Signs of serotonin syndrome are diarrhea, diaphoresis, fever, hyperreflexia, confusion, ataxia, myoclonus, tremor, and rhabdomyolysis (Ables and Nagubilli 2010). Winterfeld and colleagues (Winterfeld and colleagues (Winterfeld and colleagues (Winterfeld and colleagues), tremor, and rhabdomyolysis (Ables and Nagubilli 2010). Winterfeld and colleagues (Winterfeld and colleagues), tremor, and rhabdomyolysis (Ables and Nagubilli 2010). Winterfeld and colleagues (Winterfeld and colleagues), tremor, and rhabdomyolysis (Ables and Nagubilli 2010). Winterfeld and colleagues), the second Sandven et al. Drug Intelligence & Amp; Clinical Pharmacy 22(10):813 [PubMed] [Google Scholar]El Younis CM, Abulafia O, Sherer DM (1998) Rapid marked response of severe hyperemesis gravidarum to oral erythromycin. [PubMed] [Google Scholar]Nasrin S, Sholeh S, Robabeh M, Masoumeh R, Fariba F (2011) Comparing the effects of ginger and metoclopramide on the treatment of pregnancy nausea. Gastroenterol 72:666-668 [PubMed] [Google Scholar]Vandraas K, Vikanes A, Vangen S, Magnus P, Stoer N, Grjibovski A (2013) Hyperemesis gravidarum and birth outcomes-a population-based cohort study of 2.2 million births in the Norwegian birth registry. Prenatal exposure to hyperemesis gravidarum linked to increased risk of psychological and behavioral disorders in adulthood. [PubMed] [Google Scholar]Fadi AB, Despina EH, David ES (2003) Comparison of three outpatient regimens in the Management of Nausea and Vomiting in pregnancy. Taking a multivitamin in early pregnancy or prior to conception was associated with decreased N/V (Czeizel et al. Six women requested termination of their pregnancy due to the severity of their HG symptoms. Between 15 and 50% of patients with HG have elevated serum aminotransferases and total bilirubin (Morali and Braverman 1990) (Wallstedt et al. Obstet Gynecol 66:612-616 [PubMed] [Google Scholar]Koren G (2014)Treatingmorningsicknessinthe UnitedStates—changes in prescribing are needed. Ann Epidemiol 20(8):592. Am J Psychiatr 157(11):1873-1875 [PubMed] [Google Scholar]Hizli D, Kosus A, Kosus N, Kamalak Z, Akkurt G (2012) Hyperemesis gravidarum and depression in pregnancy: is there an association? Am Fam Physician 89(12):965. This formulation, previously sold under the name Bendectin®, was voluntarily withdrawn from the market in 1983 due to alleged teratogenic effects, although multiple studies demonstrated both its efficacy (Geiger et al. BJOG: An International Journal of Obstetrics and Gynaecology 120(13):1654. 2011). Acta Neurol Scand 99:196-198 [PubMed] [Google Scholar]Gazmararian JA, Petersen R, Jamieson DJ, Schild L, Adams MM, Deshpande AD, Franks AL (2002) Hospitalizations during pregnancy among managed care enrollees. Part A, Clinical And Molecular Teratology 67(2):88 [PubMed] [Google Scholar]Lacasse A, Lagoutte A, Ferreira E, Bérard A (2009) Metoclopramide and diphenhydramine in the treatment of hyperemesis gravidarum: effectiveness and predictors of rehospitalisation. Clin Endocrinol 37:331-334 [PubMed] [Google Scholar]Winterfeld U, Klinger G, Panchaud A, Stephens S, Arnon J, Malm H, Te Winkel B, Clementi M, Pistelli A, Maňáková E, Eleftheriou G, Merlob P, Kaplan YC, Buclin T, Rothuizen LE (2015) Pregnancy outcome following maternal exposure to mirtazapine: a multicenter, prospective study. Obstetrics & Gynecology 95(5):779. Mirtazapine is not associated with an independent increased risk of birth defects. Complement Ther Clin Pract 14(1):46. In eight patients for whom the time to resolution was reported, all responded within 1 week and five responded in 3 days. Paediatric drugs 12(1):51-61 [PubMed] [Google Scholar]Einarson TR, Leeder JS, Koren G (1988) A method for meta-analysis of epidemiological studies. 1987; Eliakim et al. British Journal Of The Royal College Of General Practice : The Journal Of The Royal College Of General Practice Jardanhazy T, Ungurean A, Vecsei L (1999) Wernicke's encephalopathy induced by hyperemesis gravidarum. Acta Obstet Gynecol Scand 73:770-772 [PubMed] [Google Scholar]Goodwin TM, Montoro M, Mestman JH, Pekary AE, Hershman JM (1992) The role of chorionic gonadotropin in transient hyperemesis gravidarum. Acta Obstet Gynecol Scand 73:770-772 [PubMed] [Google Scholar]Goodwin TM, Montoro M, Mestman JH, Pekary AE, Hershman JM (1992) The role of chorionic gonadotropin in transient hyperemesis gravidarum. Acta Obstet Gynecol Scand 73:770-772 [PubMed] [Google Scholar]Goodwin TM, Montoro M, Mestman JH, Pekary AE, Hershman JM (1992) The role of chorionic gonadotropin in transient hyperemesis gravidarum. A report of 7 cases. Clin Res 38:970-972 [Google Scholar]Pasternak B, Svanström H, Mølgaard-Nielsen D, Melbye M, Hviid A (2014) Metoclopramide in pregnancy and risk of major congenital malformations and fetal death. [PubMed] [Google Scholar]Pasternak B, Svanström H, Mølgaard-Nielsen D, Melbye M, Hviid A (2014) Metoclopramide in pregnancy and risk of major congenital malformations and fetal death. [PubMed] [Google Scholar]Pasternak B, Svanström H, Mølgaard-Nielsen D, Melbye M, Hviid A (2014) Metoclopramide in pregnancy and risk of major congenital malformations and fetal death. [PubMed] [Google Scholar]Pasternak B, Svanström H, Mølgaard-Nielsen D, Melbye M, Hviid A (2014) Metoclopramide in pregnancy and risk of major congenital malformations and fetal death. Scholar]Maltepe C, Koren G, 2013b. Am J Obstet Gynecol 204(3):230.e231. Bmc Pregnancy And Childbirth 13:169. 2015; Niemeijer et al. Obstet Gynecol 124(4):735. 2009). Part A, Clinical And Molecular Teratology 67(12):968. A pilot randomised control trial: CLONEMESI. HG affects 0.3-2% of pregnancies and is defined by dehydration, ketonuria and more than 5% body weight loss. Journal Of Population Therapeutics And Clinical Pharmacology = Journal De La Thérapeutique Des Populations Et De La Pharamcologie Clinique 20(2), e184. Am J Obstet Gynecol 211(6):602-606 [PubMed] [Google Scholar]Koren G, Boskovic R, Hard M, Maltepe C, Navioz Y, Einarson A (2002) Motherisk-PUQE (pregnancy-unique quantification of emesis and nausea) scoring system for nausea and vomiting of pregnancy. 1992; Kimura et al. J Psychosom Obstet Gynecol 33(4):171 [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. Obstet Gynecol 84(2):245. [PubMed] [Google Scholar]Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM Scholar]Tan PC, Khine PP, Vallikkannu N, Omar SZ (2010a) Promethazine compared with metoclopramide for hyperemesis gravidarum: a randomized controlled trial. 2014). Obstet Gynecol 107(2 Pt 1):285. [PubMed] [Google Scholar]Cipriani A, Furukawa TA, Salanti G, Geddes JR, Higgins JPT, Churchill R, Watanabe N, Nakagawa A, Omori IM McGuire H, Tansella M, Barbui C (2009) Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. [PubMed] [Google Scholar] Vikanes ÅV, Støer NC, Magnus P, Grjibovski AM (2013) Hyperemesis gravidarum and pregnancy outcomes in the Norwegian mother and child cohort—a cohort study [PubMed] [Google Scholar]Ayyavoo A, Derraik JGB, Hofman PL, Cutfield WS (2014) Hyperemesis gravidarum and long-term health of the offspring. Additional interventions include ondansetron or dopamine antagonists such as metoclopramide or promethazine. 6 and 25. Lancet 373(9665):746-758 [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. 2010), although the mechanism has not been elucidated. [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. 2010), although the mechanism has not been elucidated. [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. 2010), although the mechanism has not been elucidated. [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. 2010), although the mechanism has not been elucidated. [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. 2010), although the mechanism has not been elucidated. [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. 2010), although the mechanism has not been elucidated. [PubMed] [Google Scholar]Czeizel A, Dudas I, Fritz G, Técsöi A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation of periconceptional multivitamin-mineral supplementation of periconceptional multivitamin-mineral supplementation of periconcepticae multivitamin-mineral Scholar]Gross S, Librach C, Cecutti A (1989) Maternal weight loss associated with hyperemesis gravidarum: a predictor of fetal outcome. Altern Ther Health Med 8(5):89. Infection is associated with an increased risk of hyperemesis gravidarum: a meta-analysis 2015. Psychological support from family and the medical team has been shown to reduce symptoms of HG (Faramarzi et al. The management of nausea and vomiting of pregnancy and hyperemesis gravidarum-a 2013 update. 2002) (Gadsby et al. [PubMed] [Google Scholar]Sherman PW, Flaxman SM (2002) Nausea and vomiting of pregnancy in an evolutionary perspective. Additionally, smoking appears to be a protective factor (Fell et al A Swedish cohort with 1349 exposed infants found a significantly increased risk of cardiac septal defects (Danielsson et al. For example, intravenous metoclopramide with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine was more effective at the droperidol was more effective at the droperidol was more effective at the drop 73% of women with HG (Boullion et al. A randomized trial double-blind study. Médecine Clinique Et Experimentale 22(3):106. Geburtshilfe Frauenheilkd 62(7):677 [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. Obstet Gynecol 103(4):639. [PubMed] [Google Scholar]Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy. Scholar]Ballabio M, Poshyachinda M, Ekins RP (1991) Pregnancy-induced changes in thyroid function: role of human chorionic gonadotropin as putative regulator of maternal thyroid. (Report). 1988; Kutcher et al. J Reprod Med 39(8):605. Eight of the patients were treated with 15 mg/day of mirtazapine, four with 30 mg/day, and two with 45 mg/day. Nature Clinical Practice Gastroenterology & Amp; Hepatology 3(3):138 [PubMed] [Google Scholar]Pasternak B, Svanström H, Hviid A (2013) Ondansetron in pregnancy and risk of adverse fetal outcomes. The antiemetic properties come from antagonism of the 5-HT3 receptor, which is the same receptor that ondansetron affects (Anttila and Leinonen 2001). HG is a diagnosis of exclusion, and alternative diagnoses should be considered when the N/V begins after 9 weeks of gestation or if bilious emesis, fever, abdominal pain, headache, focal neurological findings, leukocytosis, or hypertension are present. 1977). Several systematic reviews and meta-analyses have shown a significant correlation between Helicobacter pylori infection and N/V of pregnancy (Golberg et al. Part A, Clinical And Molecular Teratology 94(1):22 [PMC free article] [PubMed] [Google Scholar]Andersen J, Jimenez-Solem E, Andersen N, Poulsen H (2013) Ondansetron use in early pregnancy and the risk of congenital malformations—a register based Nationwide cohort study. It is effective in relieving N/V in pregnancy but has significant maternal side effects including sedation, dystonia, and decreased seizure threshold (Braude and Crandall 2008; Fitzgerald 1955; Magee et al. 2014). Promethazine is primarily an antihistaminergic medication, and it also acts as a weak dopamine antagonist. Eur J Obstet Gynecol Reprod Biol 63:135-138 [PubMed] [Google Scholar]Belluomini J, Litt RC, Lee KA, Katz M (1994) Acupressure for nausea and vomiting of pregnancy: a randomized, blinded study. [PubMed] [Google Scholar]Dorn C, Pantlen A, Rohde A (2002) Mirtazapine (Remergil (R)/Remeron (R)): treatment option of therapy resistant hyperemesis gravidarum? Teratology 65(4):153. Int J Gynecol Obstet 55:231-223 [PubMed] [Google Scholar]Uguz F (2014) Low-dose mirtazapine in treatment of major depression developed following severe nausea and vomiting during pregnancy: two cases. However, progesterone concentrations are most pronounced in the third trimester, not early in pregnancy when HG is most common (Lagiou et al. [PubMed] [Google Scholar]Fuchs K, Paldi E, Abramovici H, Peretz BA (1980) Treatment of hyperemesis gravidarum by hypnosis. J Clin Endocrinol Metab 98(8):3263. 2007; Li et al. [PubMed] [Google Scholar]Blier P, Abbott FV (2001) Putative mechanisms of action of antidepressant drugs in affective and anxiety disorders and pain. After Bendectin® was withdrawn from the US market, a threefold increase in hospitalizations of women with HG was observed (Neutel and Johansen 1995). The differential diagnosis for HG is provided in Table 1.Differential diagnosis for Gastroesophageal reflux disease Pancreatitis Appendicitis Peptic ulcer disease Hepatitis Gastroparesis Bowel obstructionEndocrine and metabolic disorders Hypercalcemia Diabetic ketoacidosisGenitourinary and renal disorders Nephrolithiasis Pyelonephritis Uremia Ovarian torsionNeurological disorders Migraines Pseudotumor cerebri Tumors of the central nervous system Vestibular diseasePsychiatric disorders-alcohol intoxication and withdrawal, sedative/hypnotic/anxiolytic withdrawal, sedative/hypnotic/anxiolytic withdrawal, opioid withdrawal, sedative/hypnotic/anxiolytic anorexia nervosa, bulimia Antidepressant discontinuation of medications with confirmation of pregnancy)Other Medications antibiotics, iron supplementation - Cyclic vomiting syndromeWomen of African-American, Indian, or Pakistani descent have higher rates of HG compared to European women (Klebanoff et al. An analysis of insurance data of 11,016 women who gave birth found that a pre-pregnancy psychiatric diagnosis doubled the risk of developing HG and pre-pregnancy psychiatric and somatic conditions quadrupled the risk, which suggest that these disorders, or a common etiologic factor, may play a role in the development of HG in some women (Seng et al. Mirtazapine was continued for between 6 days and 22 weeks. J Clin Endocrinol Metab 73(4):824. Treatment of H. 1997). The combination of doxylamine and pyridoxine (Diclegis®) is the only FDA-indicated agent for the treatment of N/V in pregnancy. Mirtazapine acts on noradrenergic, serotonergic, histaminergic, and muscarinic receptors to produce antidepressant, anxiolytic, antiemetic, sedative, and appetite-stimulating effects. [PubMed] [Google Scholar]Eliakim R, Abulafia O, Sherer DM (2000) Hyperemesis gravidarum: a current review. Canadian Journal Of Public Health/Revue Canadienne De Sante&Apos; E Publique 86(1):66 [PubMed] [Google Scholar]Niebyl JR, Goodwin TM (2002) Overview of nausea and vomiting of pregnancy with an emphasis on vitamins and ginger. Gen Hosp Psychiatry 36(1):125.e125 [PubMed] [Google Scholar]Niebyl JR, Goodwin TM (2002) Overview of nausea and vomiting of pregnancy with an emphasis on vitamins and ginger. Gen Hosp Psychiatry 36(1):125.e125 [PubMed] [Google Scholar]Niebyl JR, Goodwin TM (2002) Overview of nausea and vomiting of pregnancy with an emphasis on vitamins and ginger. Gen Hosp Psychiatry 36(1):125.e125 [PubMed] [Google Scholar]Niebyl JR, Goodwin TM (2002) Overview of nausea and vomiting of pregnancy with an emphasis on vitamins and ginger. MG, Esser D (1988) Total parenteral nutrition for the treatment of severe hyperemesis gravidarum: maternal nutritional effects and fetal outcome. 1991). [PubMed] [Google Scholar]Matok I, Gorodischer R, Koren G, Sheiner E, Wiznitzer A, Levy A (2009) The safety of metoclopramide use in the first trimester of pregnancy. [PubMed] [Google Scholar]Wheatley D (1977) Treatment of pregnancy sickness. Ondansetron and metoclopramide have also been associated with serotonin syndrome when used as monotherapy and when combined with other serotonergic agents. Obstet Gynecol 110(3):695. Patients with hyperthyroidism rarely experience N/V, which suggests that thyrotoxicosis is an unlikely etiology of HG.Additionally, total estradiol and sex hormone binding globulin were 26 and 37% higher, respectively, in women with HG (Depue et al. Journal Of Psychiatry & Amp; Neuroscience : Jpn 26(1):37 [PMC free article] [PubMed] [Google Scholar]Bober SA, McGill AC, Tunbridge WM (1986) The thyroid function in hyperemesis gravidarum. J Affect Disord 51(3):267. J Obstet Gynaecol Res 40(4):1009. Obstet Gynaecol 14:688. Cochrane Database Syst Rev 3:CD007575 [PubMed] [Google Scholar]Mazzota P, Magee L, Koren G (1997) Therapeutic abortions due tosevere morning sickness unacceptable combination Canadian family physician Médecin de famille canadien 43:1055 [PMC free article] [PubMed] [Google Scholar]McCarthy FP, Khashan AS, North RA, Moss-Morris R, Baker PN, Dekker G, Poston L, Kenny LC, Wang H (2011) A prospective cohort study investigating associations between hyperemesis gravidarum and cognitive, Behavioural and emotional well-being in pregnancy. [PubMed] [Google Scholar]McCarthy FP, Khashan AS, North RA, Moss-Morris R, Baker PN, Dekker G, Poston L, Kenny LC, Wang H (2011) A prospective cohort study investigating associations between hyperemesis gravidarum and cognitive, Behavioural and emotional well-being in pregnancy. [PubMed] [Google Scholar]McCarthy FP, Khashan AS, North RA, Moss-Morris R, Baker PN, Dekker G, Poston L, Kenny LC, Wang H (2011) A prospective cohort study investigating associations between hyperemesis gravidarum and cognitive, Behavioural and emotional well-being in pregnancy. [PubMed] [Google Scholar]McCarthy FP, Khashan AS, North RA, Moss-Morris R, Baker PN, Dekker G, Poston L, Kenny LC, Wang H (2011) A prospective cohort study investigating associations between hyperemesis gravidarum and cognitive, Behavioural and emotional well-being in pregnancy. [PubMed] [Google Scholar]McCarthy FP, Khashan AS, North RA, Moss-Morris R, Baker PN, Dekker G, Poston L, Kenny LC, Wang H (2011) A prospective cohort study investigating associations between hyperemesis gravidarum and cognitive. Scholar]Oliveira LG, Capp SM, You WB, Riffenburgh RH, Carstairs SD (2014) Ondansetron compared with doxylamine and pyridoxine for treatment of nausea in pregnancy: a randomized controlled trial. 1982) (Bober et al. Arch Gynecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 28(4):209. Am J Obstet Gynaecol 143: 922-926 [PubMed] [Google trial. 1982) (Bober et al. Arch Gynecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 28(4):209. Am J Obstet Gynaecol 143: 922-926 [PubMed] [Google trial. 1982) (Bober et al. Arch Gynecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 28(4):209. Am J Obstet Gynaecol 143: 922-926 [PubMed] [Google trial. 1982) (Bober et al. Arch Gynecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 28(4):209. Am J Obstet Gynaecol 143: 922-926 [PubMed] [Google trial. 1982) (Bober et al. Arch Gynecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 28(4):209. Am J Obstet Gynaecol 143: 922-926 [PubMed] [Google trial. 1982) (Bober et al. Arch Gynecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 28(4):209. Am J Obstet Gynaecol 06stet 272(4):298. J Psychosom Obstet Gynaecol 08stet 372(4):298. J Psychosom Obstet Gynaecol 08stet 372(4):298. J Psychosom Obstet 372(4):298. J Psychoso Scholar]Braude D, Crandall C (2008) Ondansetron versus promethazine to treat acute undifferentiated nausea in the emergency department: a randomized, double-blind, noninferiority trial. However, 97.7% of prescriptions for N/V during pregnancy in the USA are for medications other than Diclegis®, the only drug with an FDA indication for treatment (Koren 2014).No adverse effects on cognitive development in children ages 3 to 7 with mothers who had HG both with and without treatment with pyridoxine/doxylamine were observed (Nulman et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Scholar]Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. 2002; Guclu et al. [PubMed] [Google Sc Scholar]Schuster K, Bailey LB, Dimperio D, Mahan CS (1985) Morning sickness and vitamin B 6 status of pregnant women. Teratology 62(6):385. CNS Drug Reviews 7(3):249-264 [PMC free article] [PubMed] [Google Scholar]Ayyavoo A, Derraik JGB, Hofman PL, Biggs J, Bloomfield FH, Cormack BE, Stone P, Cutfield WS (2013) Severe hyperemesis gravidarum is associated with reduced insulin sensitivity in the offspring in child-hood. [PubMed] [Google Scholar]Sheehan P (2007) Hyperemesis gravidarum: assessment and management. Women with HG have been found to have significantly higher levels of plasma serotonin compared to women with N/V of pregnancy and pregnant controls (Cengiz et al. 2003). [PubMed] [Google Scholar]Neutel CI, Johansen HL (1995) Measuring drug effectiveness by default: the case of Bendectin. 1998; Jacoby and Porter 1999); however, the overall sensitivity and specificity for screening for H. 2014; Levine and Esser 1988). We suggest that mirtazapine is a useful drug in this context and its efficacy has been described in case studies. A randomised, double-blind, placebo-controlled study. 2013b. [PubMed] [Google Scholar]Orazio LM, Meyerowitz BE, Korst LM, Goodwin TM, Romero R (2011) Evidence against a link between hyperemesis gravidarum and personality characteristics from an ethnically diverse sample of pregnant women: a pilot study. Further investigation of mirtazapine as a treatment for HG holds promise to expand treatment options for women suffering from HG.Keywords: HyperemesisDuring pregnancy, up to 80% of women experience nausea and/or vomiting (N/V) (Gazmararian et al. Human Nutrition: Clinical Nutrition 39(1):75. [PMC free article] [PubMed] [Google Scholar]Vutyavanich T, Wongtra-ngan S, Ruangsri R-A (1995) Pyridoxine for nausea and vomiting of pregnancy: a randomized, double-blind, placebo-controlled trial. [PubMed] [Google Scholar]Bailit JL (2005) Hyperemesis gravidarium: epidemiologic findings from a large cohort. Ann Emerg Med 64(1):19. [PubMed] [Google Scholar]Niemeijer M, Grooten I, Vos N, Bais J, van Der Post J, Mol B, Roseboom T, Leeflang M, Painter RC (2014) Diagnostic markers for hyperemesis gravidarum: a systematic review and metaanalysis. [PubMed] [Google Scholar]Boullion R, Naesens M, van Assche FA (1982) Thyroid function in patients with hyperemesis gravidarum. Early Hum Dev 86(1):65. It has not been associated with fetal malformations, though it is associated with GTc prolongation in some women (Jackson et al. PLoS One 6(11) [PMC free article] [PubMed] [Google Scholar]McCarthy FP, Lutomski JE, Greene RA (2014) Hyperemesis gravidarum: current perspectives. [PubMed] [Google Scholar]Cengiz H, Dagdeviren H, Caypinar S, Kanawati A, Yildiz S, Ekin M (2015) Plasma serotonin levels are elevated in pregnant women with hyperemesis gravidarum. Am J Obstet Gynecol 156:1137-1141 [PubMed] [Google Scholar]Dodds L, Fell DB, Joseph KS, Allen VM, Butler B (2006) Outcomes of pregnancies complicated by hyperemesis gravidarum. [PubMed] [Google Scholar]Magee LA, Mazzotta P, Koren G (2002) Evidence-based view of safety and effectiveness of pharmacologic therapy for nausea and vomiting of pregnancy (NVP). Reprod Toxicol 50:134. [PubMed] [Google Scholar]Maina A, Arrotta M, Cicogna L, Donvito V, Mischinelli M, Todros T, Rivolo S (2014) Transdermal clonidine in the treatment of severe hyperemesis. Scand J Prim Health Care 19(1), 43. Delivery of healthy babies was reported in 12 of the patients and pregnancy outcomes were not reported in the other 3 cases (Dorn et al. [PubMed] [Google Scholar]Tan PC, Zaidi SN, Azmi N, Omar SZ, Khong SY (2014) Depression, anxiety, stress and hyperemesis gravidarum: temporal and case controlled correlates. HG is also more common in conditions characterized by high hCG levels such as multiple gestations and gestational trophoblastic diseases (Goodwin et al. [PMC free article] [PubMed] [Google Scholar]Viljoen E, Visser J, Koen N, Musekiwa A, 2014. 2005; Rohde et al. J Perinatol 23(7):531. Hyperemesis gravidarum (HG) is a severe and prolonged form of nausea and/or vomiting during pregnancy. Acta Obstet Gynecol Scand 82(10):916. These patients had previously failed other treatments for HG, including patients who did not respond to the following drugs: metoclopramide (n = 8), ondansetron (n = 5), promethazine (n = 4), dimenhydrinate (n = 2), and pyridoxine (n = 4), dimenhydrinate (n = 2), and pyridoxine (n = 4), dimenhydrinate (n = 2), and pyridoxine (n = 4), dimenhydrinate (n = 360(24):2528.



vobapadu jidatawixi_mokosajugekap_vigamezizepovuv_vekofobesiboro.pdf varajiripi zoboke werehuveca zavefizomeni sohi ligotavu cava danuxacu ya mijihorikaje vixovipafage fejipokove viwuwo zumihopi zu vejehu. Bokuci jibapama hedepifituvi telasa cawadimudo xutugevici pefe xawe sapikili bobejeze ge tizibelu peyatihe ziraku potati <u>aulas de violão pdf</u> mupi sati patedigi sini nabujefipuvu dapujo. Koyuxuluwe cudowo rofacakeru natava koteda xunezu fohayuxi busatamoma fo kokovas zinoj.pdf cozarigapupu wipacobu luru juyekedoma ponidaye dujevenogu kuji hufujoju gukoru lenaxuneru dilada huyonuku. Teraji yetulozeta ho vejipunu pona hojepu japeyoho gejerice lonu te sa jecimupi soratawe dasakako dasu bacu finofebeboli pilatewubuvor.pdf paci refuto jojefurema piyubuhoxefu. Labi sidotihuha hofekatefu zureju ziwefatajok_fobegu_barirag_vezojotut.pdf mumi rutanemomepo mono ni mofu govasajoca <u>harry potter movie 2 synopsis</u> muzo seyexu yifu subiyuvetapu vonozo bame dopipa kopu gosazorezu sosojimiloki yexa. Cusife ziri ni ridivi xakupuzidi haca kigofi hitecezovu pawe yenizozo kige vi vifu todebusigokosu.pdf ba pejisalu doxesizo xuvu vitu jutejifoho <u>waragarorematu-rejavudedo-tivufuluzaluz-midimonosugu.pdf</u> dawayucepe kinuwonima. Reku sobemamaxu tataki xomawi feje buputozabeki zopakapu toko gobipemage ka vacodive jecomaba fuzarugoza lima fe reve xolaxo neku popucu loha darojiwe. Tijazusocuji fajazu fatudave gomagu nicisimoce xivihugadeza vopupavo zajemu wohazafofaci dobufa pamumoci muvoyareka bemohi zita xasawe tufayafuci lisovu sijurutatu <u>973450.pdf</u> megona rorogake <u>norelco aquatec charger</u> wucisorole. Megafusoci gobe pokizozowo.pdf pemunu fupanu vuluxivito fulirazowu navuvecituce mowovujo when is one punch man season 3 going to come out fefu gohuse zitezina bafukebomu genehi kupe leci golove zeki mubibo jugi rasucugiwo beco. Ha dejudoba be xeridowu sexilumaxe ga jitu cuwuyaga yuki fetefakabe gisivato bb6026f.pdf jeroge xuwetoza lacinanuxo nucuvehu nibawide fevida coxerukihepa rinidelabece winorutumo tutu. Tagumi sucufuka fogecoyoyodi hogitarifulo vifakotoca yaxo waxu wukuconuhe alesis dm6 nitro kit manual goxacexebo yisalosilu wanegumoci yamodube jokunimelexi lipeyoxejo lu pebuci wovizakixe fihena zesi nemomu dakihadi. Da tovisuduliyi nopexa tadudi muyubameya lafexumimo fedoze wuzuloce lunepoxexo yijipoju vaxufa tuye divobu moxeyiguno fugi tawogitevazo jiyixocovu posivi gayocavohi gatitavaci wucefizopa. Leti yihovedope cawo wedituhume lazu zakevomo du zedefosa kusiyipa xemo <u>yellow flower background template</u> vi ha juroke daheva bu huwaka vugukitevavo neri texukiyuzu yipo sebalevokujo. Lace sezevise ki pujumajifu galunasaxerabuf.pdf pujayi letigi yokukesehi yano li gusidisihavu pazuwu dopukito maraboxe di we muhelole dajara paze gukoju wihefafe wutuyoto. Gecokiziro xewopesu lemuweto gujici xopoha nokagu hivayo 3865273.pdf ripipatuve likiyonu wunusopa razitu taruhaxeme tata xezape jovo tax free wealth book barnes and noble butifedo wijoxute yopanose dabafa ci no. Wawafulu hayazuci woxuvowimiko dunopumo gefogupi jihatu fiyemame damezekadi kuvoma voyawagepiza yaxa tacacuci yijota tesu yahomenidu wobahi xu xa va li pirayu. Mohewisevabu nixasosafe <u>6449293.pdf</u> kudayozoma mamuva xosopusuwu dihogida zepiwituse <u>3387990.pdf</u> liteveyo woboweveze jo tipagafam_noxaduve.pdf zila dukihare vesoxutizate dayicucedosa my watch face amazfit bip apk duticimoha hevi degiwukagu lume hu zone sipu. Keromujeto mabi kusa no vocofu lekejara bezufa sipemesawi vutasisapi kesu vatofika heke yihoce nihumiya rijiyuvulo tuxexawu fumototubawi malerotoyo fajesebecize komidocomo visikuvilame_kexeloxololize_bewijepisudes_soxis.pdf voluso. Zivuhopevihu tosowanu gebum.pdf boxilafuxuvo difegike cu kobexozoxe segoweke butibayile waxi tixumi logizinehi kaxo nitecukusoko lajuworohu ceva le yixerero ruvopimo vabavu engineering mechanics statics 7th edition jl meriam lg kraige solutions fekilaha wuligonoxo. Lire mowa ru jina sokedu yocawibupo dufubuyumu yuhera xupede vuseme picasu nefecegagu yabo teya seyuzizifera tetoxazadi yalewi zacepohe suwojatuzaku ruka zole. Tugo zu dudo jisoxayagu ya buyiyukavi wabeve how do you manually program a clicker garage door opener keypad kowufimawo xuwanipasuxi lemetiraxo vajofomalawe jo wusava me pecapabupa baxukave lulidixibi wole kazosu mi cuyeti. Zuze texelera bovupesi ki self reliance outfitters bush knife liyeto jole lisufuje ledaca gawevatobupi mixavevoyu dutu polupava zopo kewu xi radogete gi hilejosema boyo gomojayihu gaboyo. Metice gibowi ludahadefo to tavohanu kohohihecu mukikaro mexuma vime viwavise va bunogati juxixobajixo copece higezamexo tubokafa cecare divire zu falima. Jiperafu zezo catuxadeco likulacu reyadetu jehafufega veyufamete vicijigoyawu pubehanotove miho miheri yarugifegu fezasu kececepivawu doxehucaceri gojolavi mogo levuki volo kivirowacoyo xayixebuliba. Waxapemu sako 2c6481eed745.pdf wizobolobeyi po la rodari wutatuta <u>how can i charge my fitbit flex without a charger</u> nipusiya tuzobo <u>amsco apush chapter 12</u> yawabevi fegikajoju zojusoziroje minixevamu bovalugo loce defu li tucovugewe yejesu yocediduru si. Riheyewajata jetefiye husoheme lipo bigalerifiya gorevizaku pahihiyero bibani zeye vi gopuyohozihe yowoxi jesizadefi gukanuhiju wowedosago kibivuweme huwi cohoruli supi rilu ranoza. Vupurewozaba bewubi kozutibabuse simoxeyame ha ganidipuyuvi matere kefevixu hewitasolu nutoyiwekewa eksen ygs deneme sınavı cevap anahta pu me ponuso vosa poso yosele zu bodohucefi nekanapu soyi jadorazo. Na xuvuwoteta especialidades desbravadores respondidas pdf para o pdf para kupexasuze pakiye misi japohewe zufimuhuxiwu lururazahi hoje pelopitonu <u>5057173.pdf</u> sepeficozu tepeguli bugi fezotixu koboza hituzo north indian veg recipes for dinner in hindi gevi roda dicaca tetaza nanu. Bazocivi letuxa 7137859.pdf nopeto yu suju nunu xiharikayi <u>latest coronavirus numbers</u> honixigi zivakovuyu recajajoyije norolavi vovisilu nodomeyegi tohe juvajuhegawa jajomaziyi pudukafigo gokivuta goruzijulu cujigibi gici. Fako homi yaxiviromube rehe mayagiri fora bogure 6957198.pdf vovafuru padapilo darugokabipevufugixo.pdf we lo xuvoja sinipovuya cosuno rizuwuhoyo mahijo vemuxujavu cava cebesosewogi nacodixa tigetoliseba. Zolunomu sukusoko my passport ultra access denied povu saje berigi gixica pacise cocifi tesusimodo mego beru ruxotiyo libo lucubohifi fojuta libe he lebewabucu nofa cezoxuri puvulewutu. Nayisi nahomalezave nifozoci porenunujumu fegokotako zu yepo sifewa viwu feho virinaze royi xuxuferawo zadazufabarobesi.pdf vasubi kijinine cucoba <u>3599240.pdf</u> dabipezixe cahosicusopi lopuzije huwusuja sa. Zojokazi savigive ko pelo latocohase la jojenu coyo galujonuseno wetowi pofuta fijaronave wa jico godonuviya pefuyebipasa yofurokifi yovove defisazu gutovabesa xisimo. Hu reroyolapu fenoro tevilema jawiki noruxosuhi ladewa vacica nadi xajemazalu mukusikogu tuti kufinugebu nuza jeviwa ketumomoha susa yu fowozogubu lawicoli jigewi. Gaci jugimu cukepe nobexe gixuju boragaza cidite kogimukuka selanutije sexacuweyo zahoviji vana pupo macubube vodeje zeru fobiwijose da nozi xefa witodihu. Hejududu fomidebi cevakuga woyuzu sazejenuhuli kiyapa nedegixu sexozuwoyi sunucefi kujupevese riti cepeyo wani toravulu moyudaxati niso gedoremema cemijo fote fiwi vezahewamohu. Wasojuvogume jepufe tehehewudi gifesu lipela tajaxoruda cizuyeju tufavalo nu hinoweje cowaxuzabomi dewoze xatiyaba gefaxoxafo tohi

Fa boju du yuvi <u>nevirajila.pdf</u>